Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 17 November 2011 (MN LaTeX style file v2.2)

Models of thin discs and spheroidal haloes with masses in a linear relationship: mass estimates for NGC4389 and UGC6969

Guillermo A. González[1]*, J. Ibáñez[2]* and Jerson I. Reina[1]*
[1]*Escuela de Física, Universidad Industrial de Santander, A. A. 678, Bucaramanga, Colombia*
[2]*Departamento de Física Teórica, Universidad del País Vasco, 48080 Bilbao, Spain*

17 November 2011

arXiv:1111.3852v1 [astro-ph.CO] 16 Nov 2011

ABSTRACT
A family of models of thin discs and spheroidal haloes with masses in a linear relationship is presented. The models are obtained by considering the gravitational potential as the superposition of two independent components, a potential generated by the thin galactic disc and a potential generated by the spheroidal halo. The models leads to an expression for the circular velocity that can be adjusted very accurately to the observed rotation curves of some specific galaxies, in such a way that the models are stable against radial and vertical perturbations. Two particular models for galaxies NGC4389 and UGC6969 are obtained by adjusting the circular velocity with data taken from the recent paper by Verheijen & Sancici (2001). The values of the halo mass, the disc mass and the total mass for these two galaxies are computed in such a way that we obtain a very narrow interval of values for these quantities. Furthermore, the values of masses here obtained are in perfect agreement with the expected order of magnitude and with the relative order of magnitude between the halo mass and the disc mass.

Key words: stellar dynamics – galaxies: kinematics and dynamics.

1 INTRODUCTION

One of the oldest and most important problems in galactic dynamics is the determination of the mass distribution based on the observations of the circular velocity or rotation curve (Pierens & Hure 2004), defined as the speed of the stars moving in the galactic plane in circular orbits around the center. Now, if we assume a particular model for the composition of the galaxy, the fit of that model with the rotation curve of a particular galaxy can, in principle, completely determine the distribution of mass. So then, the rotation curve provides the most direct method to measure the distribution of mass of a galaxy (Binney & Tremaine 2008).

Currently, the most accepted description of the composition of spiral galaxies is that a significant portion of its mass is concentrated in a thin disc, while the other contributions to the total mass of the galaxy comes from a spherical halo of dark matter, a central bulge and, perhaps, a central black hole (Binney & Tremaine 2008). Now, since all components contribute to the gravitational field of the galaxy, obtaining appropriate models that include the effects of all parts is a problem of great difficulty. However, the contribution of each part is limited to certain distance scales, so in a reasonably realistic model is not necessary to include the contribution of all components (Faber 2006).

In particular, the gravitational influence of the central black hole is appreciable only within a few parsecs around the center of the galaxy (Schödel et al 2002), so it can be completely neglected when studying the dynamics in the disc, or in regions outside the central bulge, while the bulge mainly dominates the inner region of the galaxy to a few kiloparsec. So then, the main contributions to the gravitational field of the galaxy come from the galactic disc and the dark matter halo (Faber 2006). However, it is commonly accepted that many aspects of galactic dynamics can be described, in a fairly approximate way, using models that consider only the contribution of a thin galactic disc (Binney & Tremaine 2008).

Accordingly, the study of the gravitational potential generated by an idealized thin disc is a problem of great astrophysical relevance and so, through the years, different approaches have been used to obtain such kind of thin disc models (see Binney & Tremaine (2008) and references therein). So, once an expression for the gravitational potential has been derived, corresponding expressions for the surface mass density of the disc and for the circular velocity of the disc particles can be obtained. Then, if the expression for the circular velocity can be adjusted to fit the observational data of the rotation curve of a particular galaxy, the total mass can be obtained by integrating the corresponding surface mass density.

However, although most of these thin disc models have surface densities and rotation curves with remarkable properties, many of them mainly represent discs of infinite extension and thus they are rather poor flat galaxy models. Therefore, in order to obtain more realistic models of flat galaxies, is better to consider methods that permit obtaining finite thin disc models. Now, a simple method to obtain the gravitational potential, the surface density and the rota-

---

* E-mail: guillego@uis.edu.co (GAG); j.ibanez@ehu.es (JI); jerson-reina@gmail.com (JIR)

© 0000 RAS

tion curve of thin discs of finite radius was developed by Hunter (1963), the simplest example of a disc obtained by this method being the well known Kalnajs (1972) disc.

In a previous paper (González & Reina 2006) we use the Hunter method in order to obtain an infinite family of thin discs of finite radius with a well behaved surface mass density. This family of disc models was derived by requiring that the surface density behaves as a monotonously decreasing function of the radius, with a maximum at the center of the disc and vanishing at the edge. Furthermore, the motion of test particles in the gravitational fields generated by the first four members of this family was studied in Ramos-Caro, López-Suspez &González (2008). So, although the mass distribution of this family of discs presents a satisfactory behaviour in such a way that they could be considered adequate as flat galaxy models, their corresponding rotation curves do not present a so good behavior as they do not reproduce the flat region of the observed rotation curve.

On the other hand, in Pedraza, Ramos-Caro &González (2008) a new family of discs was obtained as a superposition of members of the previously obtained family, by requiring that the surface density be expressed as a well behaved function of the gravitational potential, in such a way that the corresponding distribution functions can be easily obtained. Furthermore, besides presenting a well-behaved surface density, the models also presented rotation curves with a better behavior than the generalized Kalnajs discs. However, although these discs are stable against small radial perturbations of disc star orbits, they are unstable to small perturbations vertical to the disc plane. Then, apart of the stability problems, these discs can be considered as quite adequate models in order to describe satisfactorily a great variety of galaxies.

Based on these works, in González, Plata-Plata & Ramos-Caro (2010) were obtained some thin disc models in which the circular velocities were adjusted to very accurately fit the observed rotation curves of four spiral galaxies of the Ursa Major cluster, galaxies NGC3877, NGC3917, NGC3949 and NGC4010. These models presented well behaved surface densities and the obtained values for the corresponding total mass it agrees with the expected order of magnitude. However, the models presented a central region with strong instability to small vertical perturbations. Now, this result was expected as a consequence of the fact that the models only consider the thin galactic disc. Therefore, more realistic models must be considered that include the non-thin character of the galactic disc or the mass contribution of the spheroidal halo.

In agreement with the above considerations, in this paper we will consider a family of models obtained by expressing the gravitational potential as the superposition of a potential generated by the thin galactic disc and a potential generated by the spheroidal halo, in such a way that the model implies a linear relationship between the masses of the thin disc and the spheroidal halo. By adjusting the corresponding expression for the circular velocity to the observed data of the rotation curve of some specific galaxies, some particular models will be analysed. Then, from the corresponding expressions for the disc surface density and the density of the halo, estimate values for the total mass of the disc and the total mass of the halo will be obtained. The paper is organised as follows. First, in Section 2, we present the thin disc plus halo model. Then, in Section 3, we obtain the corresponding expressions for particular models and, in Section 4, the models are fitted to data of the observed rotation curve of some galaxies of the Ursa Mayor Cluster, as reported in Verheijen & Sancici (2001). Finally, in Section 5, we discuss the obtained results.

2 THE THIN DISC PLUS HALO MODEL

In order to obtain galaxy models consisting of a thin galactic disc and a spheroidal halo, we begin considering an axially symmetric gravitational potential $\Phi = \Phi(R, z)$, where (R, φ, z) are the usual cylindrical coordinates. Also, besides the axial symmetry, we suppose that the potential has symmetry of reflection with respect to the plane $z = 0$,

$$\Phi(R, z) = \Phi(R, -z), \tag{1}$$

which implies that the normal derivative of the potential satisfies the relation

$$\frac{\partial \Phi}{\partial z}(R, -z) = -\frac{\partial \Phi}{\partial z}(R, z), \tag{2}$$

in agreement with the attractive character of the gravitational field. We also assume that $\partial\Phi/\partial z$ does not vanish on the plane $z = 0$, in order to have a thin distribution of matter that represents the disc.

On the other hand, in order to separately describe the thin disc and the spheroidal halo, we consider that the gravitational potential can be written as the superposition of two independent components

$$\Phi(R, z) = \Phi_d(R, z) + \Phi_h(R, z), \tag{3}$$

where $\Phi_d(R, z)$ is the part of the potential generated by the thin galactic disc, while $\Phi_h(R, z)$ corresponds to the spheroidal halo component. The disc component $\Phi_d(R, z)$ must be a solution of the Laplace equation everywhere outside the disc,

$$\nabla^2 \Phi_d = 0, \tag{4}$$

while the halo component $\Phi_h(R, z)$ satisfies the Poisson equation

$$\nabla^2 \Phi_h = 4\pi G \varrho, \tag{5}$$

where $\varrho(R, z)$ is the mass density of the halo.

So, given a potential $\Phi(R, z)$ with the previous properties, we can easily obtain the circular velocity $v_c(R)$, defined as the velocity of the stars moving at the galactic disc in circular orbits around the center, through the relationship

$$v_c^2(R) = R \left.\frac{\partial \Phi}{\partial R}\right|_{z=0}, \tag{6}$$

while the surface mass density $\Sigma(R)$ of the thin galactic disc is given by

$$\Sigma(R) = \frac{1}{2\pi G} \left.\frac{\partial \Phi}{\partial z}\right|_{z=0^+}, \tag{7}$$

which it is obtained by using the Gauss law and the reflection symmetry of $\Phi(R, z)$.

Accordingly, in order that the potential of the spheroidal halo does not contribute to the disc surface density, we will impose the condition

$$\left.\frac{\partial \Phi_h}{\partial z}\right|_{z=0^+} = 0. \tag{8}$$

Furthermore, in order to have a surface density corresponding to a finite disclike distribution of matter, we impose boundary conditions in the form

$$\left.\frac{\partial \Phi_d}{\partial z}\right|_{z=0^+} \neq 0; \quad R \leqslant a, \tag{9a}$$

$$\left.\frac{\partial \Phi_d}{\partial z}\right|_{z=0^+} = 0; \quad R > a, \tag{9b}$$

in such a way that the matter distribution is restricted to the disc $z = 0$, $0 \leqslant R \leqslant a$, where a is the radius of the disc.

In order to properly pose the boundary value problem, we introduce the oblate spheroidal coordinates, whose symmetry adapts in a natural way to the geometry of the model. These coordinates are related to the usual cylindrical coordinates by the relation (Morse & Fesbach 1953)

$$R = a\sqrt{(1+\xi^2)(1-\eta^2)}, \tag{10a}$$

$$z = a\xi\eta, \tag{10b}$$

where $0 \leqslant \xi < \infty$ and $-1 \leqslant \eta < 1$. The disc has the coordinates $\xi = 0$, $0 \leqslant \eta^2 < 1$. On crossing the disc, the η coordinate changes sign but does not change in absolute value. The singular behaviour of this coordinate implies that an even function of η is a continuous function everywhere but has a discontinuous η derivative at the disc.

Now, in terms of the oblate spheroidal coordinates, the Laplace operator acting over any axially symmetric function $\Phi(\xi, \eta)$ gives

$$\nabla^2\Phi = \frac{\left[(1+\xi^2)\Phi_{,\xi}\right]_{,\xi} + \left[(1-\eta^2)\Phi_{,\eta}\right]_{,\eta}}{a^2(\xi^2+\eta^2)}, \tag{11}$$

whereas the boundary condition (8) is equivalent to

$$\left.\frac{\partial\Phi_h}{\partial\xi}\right|_{\xi=0} = 0, \tag{12a}$$

$$\left.\frac{\partial\Phi_h}{\partial\eta}\right|_{\eta=0} = 0, \tag{12b}$$

and the boundary conditions (9a) and (9b) reduce to

$$\left.\frac{\partial\Phi_d}{\partial\xi}\right|_{\xi=0} \neq 0, \tag{13a}$$

$$\left.\frac{\partial\Phi_d}{\partial\eta}\right|_{\eta=0} = 0. \tag{13b}$$

Moreover, in order that the gravitational potential be continuous everywhere, $\Phi(\xi, \eta)$ must be an even function of η, which grants also the fulfilment of conditions (12b) and (13b).

Accordingly, by imposing the previous boundary conditions over the general solution of the Laplace equation in oblate spheroidal coordinates, we can write the gravitational potential of the galactic disc as (Bateman 1944)

$$\Phi_n(\xi, \eta) = -\sum_{l=0}^{n} C_{2l}\, q_{2l}(\xi) P_{2l}(\eta), \tag{14}$$

where n is a positive integer, which it defines the model of disc considered. Here $P_{2l}(\eta)$ are the usual Legendre polynomials and $q_{2l}(\xi) = i^{2l+1} Q_{2l}(i\xi)$, with $Q_{2l}(x)$ the Legendre functions of second kind (see Arfken & Weber 2005 and, for the Legendre functions of imaginary argument, Morse & Fesbach 1953, page 1328). The coefficients C_{2l} are, in principle, arbitrary constants, but which must be specified to obtain any particular model. We will do this after, by adjusting the circular velocity of the model with the observed data of the rotation curve of some specific galaxies.

With this expression for the gravitational potential of the disc, the surface density is given by

$$\Sigma(\widetilde{R}) = \frac{1}{2\pi a G \eta} \sum_{l=0}^{n} C_{2l}(2l+1) q_{2l+1}(0) P_{2l}(\eta), \tag{15}$$

where, as $\xi = 0$, $\eta = \sqrt{1 - \widetilde{R}^2}$, with $\widetilde{R} = R/a$. Then, by integrating on the total area of the disc, we find the value

$$\frac{\mathcal{M}_d G}{a} = C_0 \tag{16}$$

for the total mass of the disc. Now, it is clear that the surface density diverges at the disc edge, when $\eta = 0$, unless that we impose the condition (Hunter 1963)

$$\sum_{l=0}^{n} C_{2l}(2l+1) q_{2l+1}(0) P_{2l}(0) = 0, \tag{17}$$

that, after using the identities

$$P_{2n}(0) = (-1)^n \frac{(2n-1)!!}{(2n)!!}, \tag{18a}$$

$$q_{2n+1}(0) = \frac{(2n)!!}{(2n+1)!!}, \tag{18b}$$

which are easily obtained from the properties of the Legendre functions, leads to the expression

$$C_0 = \sum_{l=1}^{n} (-1)^{l+1} C_{2l}, \tag{19}$$

which gives, through (16), the value of the dics mass $\mathcal{M}_d$ in terms of the constants C_{2l}, with $l \geqslant 1$.

Now, to properly choose the gravitational potential of the spheroidal halo, first we consider the product functions

$$\Phi_j^k(\xi, \eta) = q_j^k(\xi) P_j^k(\eta), \tag{20}$$

where (Lamb 1945)

$$q_j^k(\xi) = (1+\xi^2)^{\frac{k}{2}} \frac{d^k q_j(\xi)}{d\xi^n}, \tag{21}$$

are the solutions of the differential equation

$$\frac{d}{d\xi}\left[(1+\xi^2)\frac{dq_j^k}{d\xi}\right] = \left[j(j+1) - \frac{k^2}{1+\xi^2}\right] q_j^k(\xi), \tag{22}$$

while the associated Legendre functions (Arfken & Weber 2005),

$$P_j^k(\eta) = (1-\eta^2)^{\frac{k}{2}} \frac{d^k P_j(\eta)}{d\eta^k}, \tag{23}$$

are the solutions of the differential equation

$$\frac{d}{d\eta}\left[(1-\eta^2)\frac{dP_j^k}{d\eta}\right] = \left[\frac{k^2}{1-\eta^2} - j(j+1)\right] P_j^k(\eta), \tag{24}$$

where j and k are integers, with $j \geqslant k$.

With the previous expressions, and using the Laplace operator (11), we can now compute the mass density associated with a gravitational potential given by $\Phi_j^k(\xi, \eta)$. Thus, from the Poisson equation (5), we obtain

$$4\pi G \varrho_j^k(\xi, \eta) = \frac{k^2 \Phi_j^k(\xi, \eta)}{a^2(1+\xi^2)(1-\eta^2)}, \tag{25}$$

with $k \neq 0$, while the density is identically zero when $k = 0$. Then, as $P_j^k(\eta)$ is finite at the interval $-1 \leqslant \eta \leqslant 1$ and $q_j^k(\xi)$ goes to zero when $\xi \to \infty$, $\varrho_j^k(\xi, \eta)$ properly vanishes at infinity. On the other hand, from (23) it is easy to see that $\varrho_j^k(\xi, \eta)$ diverges at $\eta = \pm 1$ (the z axis) unless that $k \geqslant 2$, while for $k > 2$ this function vanishes at $\eta = \pm 1$. On the other hand, due to the discontinuous character of η, $\Phi_j^k(\xi, \eta)$ will be continuous across the disc only if $(j - k)$ is an even number, since then $P_j^k(\eta)$ will be an even function of η.

In agreement with the above considerations, we will take for the gravitational potential of the spheroidal a simple combination of the product functions $\Phi_j^k(\xi, \eta)$ given by

$$\Phi_h(\xi, \eta) = B_{00}\Phi_0^0(\xi, \eta) + B_{20}\Phi_2^0(\xi, \eta) + B_{22}\Phi_2^2(\xi, \eta), \tag{26}$$

where B_{00}, B_{20} and B_{22} are, in principle, arbitrary constants. Then, after using the explicit expressions for $q_j^k(\xi)$ and $P_j^k(\eta)$, the halo density can be written as

$$\varrho(\xi) = \frac{3B_{22}}{\pi G a^2}\left\{3\cot^{-1}\xi - \frac{\xi(3\xi^2+5)}{(1+\xi^2)^2}\right\}, \tag{27}$$

which is maximum at the disc surface, when $\xi = 0$, and then fastly decreases being constant at the oblate spheroids defined by $\xi = \mathrm{cte}$. By integrating over all the space, we obtain the expression

$$B_{22} = \frac{\mathcal{M}_h G}{16a}, \tag{28}$$

where $\mathcal{M}_h$ is the total mass of the halo. Finally, from the condition (12a), we obtain

$$B_{00} = -\frac{\mathcal{M}_h G}{a}, \tag{29a}$$

$$B_{20} = \frac{\mathcal{M}_h G}{2a}, \tag{29b}$$

and so all the constants in (26) are expressed in terms of the halo mass $\mathcal{M}_h$.

On the other hand, if we restrict to particles moving in the thin disc, the circular velocity (6) is written in terms of the spheroidal coordinates as

$$v_c^2 = \frac{(\eta^2-1)}{\eta}\frac{\partial\Phi}{\partial\eta}\bigg|_{\xi=0}, \tag{30}$$

which, by using (3), (14), (26) and the properties of the Legendre functions, reduces to

$$v_c^2(\widetilde{R}) = \frac{\widetilde{R}^2}{\eta}\sum_{l=1}^{n}\widetilde{C}_{2l}P'_{2l}(\eta), \tag{31}$$

where

$$\widetilde{C}_2 = q_2(0)\left[C_2 + \frac{\mathcal{M}_h G}{4a}\right], \tag{32}$$

and

$$\widetilde{C}_{2l} = q_{2l}(0)C_{2l}, \tag{33}$$

for $l \geqslant 2$. Then, by using (19), (32) and (33), it is easy to see that

$$\frac{\mathcal{M}_d G}{a} + \frac{\mathcal{M}_h G}{4a} = \sum_{l=1}^{n}\frac{(-1)^{l+1}\widetilde{C}_{2l}}{q_{2l}(0)}, \tag{34}$$

and thus the model implies a linear relationship between $\mathcal{M}_d$ and $\mathcal{M}_h$, in which the independent term is determined by the constants $\widetilde{C}_{2l}$, with $l \geqslant 1$. Now, it is clear that the above relationship makes sense only if the right hand side it is positive, which should be checked for every set of constants $\widetilde{C}_{2l}$ corresponding to any particular model.

3 OBTAINING OF PARTICULAR MODELS

In order to obtain particular models, we must to specify the constants $\widetilde{C}_{2l}$ of the general model. So, we will adjust these constants in such a way that the circular velocity $v_c^2(\widetilde{R})$ it fits the data of the rotation curve of some particular galaxy. Now, as expression (31) for the circular velocity only involves derivatives of the Legendre polynomials of even order, it is equivalent to

$$v_c^2(\widetilde{R}) = \sum_{l=1}^{n}A_{2l}\widetilde{R}^{2l}, \tag{35}$$

where the A_{2l} constants are related with the previous constants $\widetilde{C}_{2l}$, for $l \neq 0$, through the relation

$$\widetilde{C}_{2l} = \frac{4l+1}{4l(2l+1)}\sum_{k=1}^{n}A_{2k}I_{kl}, \tag{36}$$

where

$$I_{kl} = \int_{-1}^{1}\eta(1-\eta^2)^k P'_{2l}(\eta)d\eta, \tag{37}$$

which is obtained by equaling expressions (31) and (35) and by using the orthogonality properties of the associated Legendre functions (Arfken & Weber 2005).

Then, if the constants A_{2l} are determined by a fitting of the observational data of the corresponding rotation curve, the corresponding values of the coefficients $\widetilde{C}_{2l}$ can be determined by means of relation (36), obtaining then a particular case of (34) corresponding to a specific galaxy model, which can be written in terms of the constants A_{2l} as

$$\frac{\mathcal{M}_d G}{a} + \frac{\mathcal{M}_h G}{4a} = \sum_{l=1}^{n}\sum_{k=1}^{n}\frac{(-1)^{l+1}(4l+1)A_{2k}I_{kl}}{4l(2l+1)q_{2l}(0)}; \tag{38}$$

however, this relation does not determine completely the values of $\mathcal{M}_d$ and $\mathcal{M}_h$, only gives a linear relationship between them. So, in order to restrict the allowed values of these masses, it is needed to analyse the behaviour of some other quantities characterizing the kinematics of the model.

Besides the circular velocity, two other important quantities concerning the kinematics of the models are the epicycle or radial frequency, $\kappa^2(R)$, and the vertical frequency, $\nu^2(R)$, which describe the stability against radial and vertical perturbations of particles in quasi-circular orbits (Binney & Tremaine 2008, Section 2.2.3, page 164). These two quantities, which must be positive in order to have stable circular orbits, are defined as

$$\kappa^2(R) = \frac{\partial^2\Phi_{\rm eff}}{\partial R^2}\bigg|_{z=0}, \tag{39}$$

$$\nu^2(R) = \frac{\partial^2\Phi_{\rm eff}}{\partial z^2}\bigg|_{z=0}, \tag{40}$$

where

$$\Phi_{\rm eff} = \Phi(R,z) + \frac{\ell^2}{2R^2}, \tag{41}$$

is the effective potential and $\ell = Rv_c$ is the specific axial angular momentum. Then, by using expression (6) for the circular velocity, we can write the above expressions as

$$\kappa^2(R) = \frac{1}{R}\frac{dv_c^2}{dR} + \frac{2v_c^2}{R^2}, \tag{42}$$

$$\nu^2(R) = \nabla^2\Phi\big|_{z=0} - \frac{1}{R}\frac{dv_c^2}{dR}, \tag{43}$$

where we also used the expression for the Laplace operator in cylindrical coordinates.

Now, by using (35), the epicycle frequency can be cast as

$$\widetilde{\kappa}^2(\widetilde{R}) = \sum_{l=1}^{n}2(l+1)A_{2l}\widetilde{R}^{2l-2}, \tag{44}$$

where $\widetilde{\kappa} = a\kappa$. So, as the epicycle frequency is completely determined by the set of constants A_{2l}, which are fixed by the numerical fit of the rotation curve data, there are not free parameters that can be adjusted by requiring radial stability. However, relation (44) can be used as a consistency criterion to the numerical fit of the circular velocity with the observed data of the rotation curve. So, if the epicycle frequency it is not positive everywhere at the interval $0 \leqslant \widetilde{R} \leqslant 1$ for a given set of constants, it is possible to make a new numerical fit by considering a different model, with another value of n, in order to ensure the positiveness of the epicycle frequency. Accordingly, we seek for a balance between the best numerical fit and the numerical fit that agrees with the positiveness of the epicycle frequency.

Also, by using the Poisson equation (5), the expression (27) for the halo density and the expression (35) for the circular velocity, the vertical frequency can be written as

$$\widetilde{\nu}^2(\widetilde{R}) = \frac{9\pi G\mathcal{M}_h}{8a} - f_1(\widetilde{R}), \tag{45}$$

where $\widetilde{\nu} = a\nu$ and

$$f_1(\widetilde{R}) = \sum_{l=1}^{n} 2lA_{2l}\widetilde{R}^{2l-2}. \tag{46}$$

So, as $\widetilde{\nu}^2$ must be positive everywhere at the interval $0 \leqslant \widetilde{R} \leqslant 1$ in order to have vertically stable models, expression (45) defines a lower bound for the halo mass,

$$\mathcal{M}_h \geqslant \mathcal{M}_h^- = \frac{8a}{9\pi G}\left\{\max f_1(\widetilde{R})\right\}, \tag{47}$$

that combined with (38) gives also an upper bound for the disc mass $\mathcal{M}_d$.

We also need to consider the behavior of the surface mass density, as given by expression (15), that can be written in terms of the constants A_{2l} as

$$\Sigma(\widetilde{R}) = \frac{\sqrt{1-\widetilde{R}^2}}{\pi^2 aG}\left\{f_2(\widetilde{R}) - \frac{3\pi G\mathcal{M}_h}{8a}\right\}, \tag{48}$$

where, for the model with $n = 4$,

$$\begin{aligned} f_2(\widetilde{R}) \;\; = \;\; & \left[2A_2 + \frac{8A_4}{9} + \frac{16A_6}{25} + \frac{128A_8}{245}\right] \\ & + \left[\frac{16A_4}{9} + \frac{64A_6}{75} + \frac{768A_8}{1225}\right]\widetilde{R}^2 \\ & + \left[\frac{128A_6}{75} + \frac{1024A_8}{1225}\right]\widetilde{R}^4 + \left[\frac{2048A_8}{1225}\right]\widetilde{R}^6, \end{aligned} \tag{49}$$

and for any other value of n the explicit expression of $f_2(\widetilde{R})$ can be easily obtained from (15), (32), (33) and (36). Accordingly, in order that the surface mass density be positive everywhere on the disc, expression (48) defines an upper bound for the halo mass,

$$\mathcal{M}_h \leqslant \mathcal{M}_h^+ = \frac{8a}{3\pi G}\left\{\min f_2(\widetilde{R})\right\}, \tag{50}$$

that combined with (38) gives also a lower bound for the disc mass $\mathcal{M}_d$. However, in order that the models make sense, we must have that

$$\mathcal{M}_h^+ \geqslant \mathcal{M}_h^-, \tag{51}$$

which should be checked for every set of constants corresponding to any particular model.

4 FITTING OF DATA TO THE MODELS

In order to check the applicability of the previous model to describe real galaxies, we analyse a set data taken from the recent paper by Verheijen & Sancici (2001), in which are reported the results of an extensive 21 cm-line synthesis imaging survey of 43 galaxies in the nearby Ursa Major cluster using the Westerbork Synthesis Radio Telescope. For each rotation curve data, we take the value of a, the radius of the galactic disc, as given by the last tabulated value of the radius. Accordingly, we are assuming that the radius of the galaxies is defined by the farthest observed data. Now, although this assumption about the galactic radius do not agrees with the accepted standard about the edge of the stellar disc (Binney & Merrifield 1998), we will make it since we are assuming that all the stars moving in circular orbits at the galactic plane are inside the disc and that there are no stars moving outside the disc.

Table 1. Galactic radius a (kpc) and constants A_{2l} $(\mathrm{km^2 s^{-2}})$.

	a	A_2	A_4	A_6	A_8
NGC4389	5.5	29495.4	−53607.4	60170.1	−23962.6
UGC6969	4.6	15094.2	−38417.9	56183.2	−26619.2

Table 2. Halo mass $\mathcal{M}_h$, disc mass, $\mathcal{M}_d$ and total mass $\mathcal{M}$ (kg).

	$\mathcal{M}_h$ (10^{34} kg)	$\mathcal{M}_d$ (10^{33} kg)	$\mathcal{M}$ (10^{34} kg)
NGC4389	4.24 − 4.52	3.01 − 3.70	4.62 − 4.82
UGC6969	1.81 − 1.83	1.42 − 1.47	1.96 − 1.98

Table 3. Halo mass $\mathcal{M}_h$, disc mass, $\mathcal{M}_d$ and total mass $\mathcal{M}$ $(\mathcal{M}_\odot)$.

	$\mathcal{M}_h$ $(\mathcal{M}_\odot)$	$\mathcal{M}_d$ $(\mathcal{M}_\odot)$	$\mathcal{M}$ $(\mathcal{M}_\odot)$
NGC4389	21345 − 22738	1514 − 1862	23207 − 24252
UGC6969	9116 − 9219	712 − 738	9854 − 9931

So, by taking the radius normalized in units of a, we make a non-linear least square fit of the data with the general relation (35) for all the galaxies reported by Verheijen & Sancici (2001), seeking for a balance between the best numerical fit and the fit that agrees with the positiveness of the epicycle frequency by using the consistency criterion (44) over the constants A_{2l}. By selecting between all the galaxies only those that fulfill with the consistency criterion, then we proceed to determine the lower and upper bounds over the halo mass by using expressions (47) and (50). Finally, we check that this two values of the halo mass are in agreement with the condition (51) in order that the models make sense. However, when we check the consistency of the adjust, we found that only the fit of the data for the galaxies NGC4389 and UGC6969 it agrees with this conditions, whereas that for all the other galaxies we found that $\mathcal{M}_n^+ < \mathcal{M}_n^-$.

In Table 1 we present the values of the galactic radius a and the constants A_{2l} obtained by the numerical adjust with the rotation curve data for galaxies NGC4389 and UGC6969 by taking a model with $n = 4$. With this values for the constants, we obtain for the galaxy NGC4389 the relationship

$$\mathcal{M}_h + 4\mathcal{M}_d = 5.7255 \times 10^{34} \text{ kg}, \tag{52}$$

and for the galaxy UGC6969 the relationship

$$\mathcal{M}_h + 4\mathcal{M}_d = 2.3995 \times 10^{34} \text{ kg}. \tag{53}$$

We also compute the minimum and maximum values of the halo mass $\mathcal{M}_h$, the disc mass $\mathcal{M}_d$ and the total mass $\mathcal{M}$, defined as

$$\mathcal{M} = \mathcal{M}_h + \mathcal{M}_d. \tag{54}$$



Figure 1. Plots of the circular velocity v_c in $\mathrm{km/s}$, as functions of the dimensionless radial coordinate $\widetilde{R} = R/a$, for the galaxies NGC4389 and UGC6969.



Figure 2. Plots of the epicycle frequency $\widetilde{\kappa}^2 \times 10^{-3}$ in $(\mathrm{km/s})^2$, as functions of the dimensionless radial coordinate $\widetilde{R} = R/a$, for the galaxies NGC4389 and UGC6969.

So, in Table 2, we present the values for the halo mass, the disc mass and the total mass in kg, whereas that in Table 3 we present the corresponding values in solar mass units $(\mathcal{M}_\odot)$.

In Fig. 1, we show the adjusted rotation curve for these two galaxies. The upper plot corresponds to the model for galaxy NGC4389, while the lower plot to the model for galaxy UGC6969. The points with error bars are the observations as reported in Verheijen & Sancici (2001), while the solid line are the rotation curves determined from (35) with $n = 4$, and the A_{2l} given by the numerical fit. As we can see, with the two galaxies we get a fairly accurate numerical adjustment to the rotation curve. We also plot, in Fig. 2, the epicycle frequency for the two galaxies, and we can see that this quantity is always positive in both of them.

In Fig. 3 we present the vertical frequency for the two models, the model for galaxy NGC4389 in the upper plot and the model for galaxy UGC6969 in the lower plot. In both of the models we plot the vertical frequency for the minimum value $\mathcal{M}_h^-$ of the halo mass (upper curve of each plot) and the maximum value $\mathcal{M}_h^+$ of the halo mass (lower curve of each plot). As we can see, for the two galaxies we obtain a positive vertical frequency for all the disc extension, so the models are stable against vertical perturbations. Now then, for galaxy UGC6969 the two curves are almost coincident given that the interval between the two mass values is very narrow.

Finally, in Fig. 4, we present the corresponding plots of the surface mass density for the two galaxies, again with the model for galaxy NGC4389 in the upper plot and the model for galaxy UGC6969 in the lower plot, plotting the surface mass density for the same two values of the halo mass as in the plots of the vertical frequency. For galaxy NGC4389, the density has a maximum at the disc centre and then decreases as $\widetilde{R}$ increases, vanishing at the disc edge. On the other hand, for galaxy UGC6969 the density behaves in a similar way, having a maximum at the disc centre and then decreasing, presenting then a small region where the density remains almost constant and then decreasing again to vanish at the disc edge. Furthermore, as with the vertical frequency, for galaxy UGC6969 the two curves are almost coincident again as a consequence of the narrow interval between the two values of the halo mass.

5 DISCUSSION

We presented a family of galactic models characterised by a linear relationship between the halo mass and the disc mass. The models are stable against radial and vertical perturbations and their circular velocities can be adjusted very accurately to the observed rotation curves of some specific galaxies. The here presented models are a generalisation of the models presented in González, Plata-Plata & Ramos-Caro (2010), where we only considered models with a thin galactic disc. The generalisation was obtained by adding to the gravitational potential of the thin disc the gravitational potential corresponding to a spheroidal halo, in such a way that we solve the problem of vertical unstability presented by the previous models.


NGC4389: Vertical Frequency

UGC6969: Vertical Frequency

Figure 3. Plots of the vertical frequency $\widetilde{\nu}^2 \times 10^{-3}$ in $(\mathrm{km/s})^2$, as functions of the dimensionless radial coordinate $\widetilde{R} = R/a$, for the galaxies NGC4389 and UGC6969.


NGC4389: Surface Mass Density

UGC6969: Surface Mass Density

Figure 4. Plots of the surface mass density $\Sigma \times 10^{-2}$ in $(\mathrm{kg/m^2})$, as functions of the dimensionless radial coordinate $\widetilde{R} = R/a$, for the galaxies NGC4389 and UGC6969.

Two particular models were obtained by a numerical fit of the general expression (35) for the circular velocity with the observed data of the rotation curve of galaxies NGC4389 and UGC6969. For these two galaxies we obtained a fairly accurate numerical adjustment with the rotation curve and, from the constants A_{2l} obtained with the numerical fit, we compute the values of the halo mass, the disc mass and the total mass for these two galaxies in such a way that we obtain a very narrow interval of values for these quantities. Furthermore, the values of masses here obtained are in perfect agreement with the expected order of magnitude and with the relative order of magnitude between the halo mass and the disc mass.

However, although we tested the applicability of the present model with all the galaxies reported by Verheijen & Sancici (2001), consistent models were obtained only for the two galaxies NGC4389 and UGC6969, whereas for all the other galaxies were obtained models with values of the halo mass such that $\mathcal{M}_n^+ < \mathcal{M}_n^-$. Now, this negative result can be considered as due to the simple halo model that we have taken here. Indeed, as we can see from expressions (25) and (26), only one term of the gravitational potential of the halo contributes to their density, what leaves only one free constant to be determined in order to fit the model to the imposed consistency conditions. This constant is precisely the mass of the halo, $\mathcal{M}_h$, which is determined by requiring the positiveness of the vertical frequency and the surface mass density. On the other hand, by considering additional terms in expression (26) for the halo potential, we will have new free parameters that perhaps allow to adjust the model to properly describe the behavior of other galaxies besides the two considered here.

In agreement with the above considerations, we can consider the simple set of models here presented as a fairly good approximation to obtaining quite realistic models of galaxies. In particular, we believe that the values of mass obtained for the two galaxies here studied may be taken as a very accurate estimate of the upper and lower bounds for the mass of the galactic disc and for the mass of the spheroidal halo in these two galaxies. Accordingly, we are now working on a more involved model, obtained by including additional terms in expression (26) for the halo potential, in order to get some particular models that can be properly adjusted with the observed data of the rotation curve of some other galaxies besides the two here considered.

ACKNOWLEDGMENTS

GAG and JIR want to thank Vicerrectoría de Investigación y Extensión, Universidad Industrial de Santander, by the financial support. JI thanks support from the Spanish Ministry of Science and Innovation (grant FIS2010-15492). Also GAG is grateful for the warm hospitality of Departamento de Física Teórica, Universidad del País Vasco, where a main part of this work was made.

REFERENCES

Arfken G., Weber H., 2005, Mathematical Methods for Physicists. 6th Ed. Academic Press.

Bateman H., 1944, Partial Differential Equations. Dover.

Binney J., Merrifield M., 1998, Galactic Astronomy. Princeton University Press.

Binney J., Tremaine S., 2008, Galactic Dynamics. 2nd ed. Princeton University Press.

Faber T., 2006, *Galactic halos and gravastars: static spherically symmetric spacetimes in modern general relativity and astrophysics*, M. Sc. Thesis in Applied Mathematics, Victoria University of Wellington.

González G., Plata-Plata S., Ramos-Caro J., 2010, MNRAS, 404, 468.

González G. A., Reina J. I., 2006, MNRAS 371, 1873.

Hunter C., 1963, MNRAS, 126, 299

Kalnajs A. J., 1972, ApJ, 175, 63.

Lamb H., 1945, Hydrodynamics. 6th Ed. Dover.

Morse P. M., Fesbach H., 1953, Methods of Theoretical Physics. Mc Graw Hill, New York.

Pedraza J. F., Ramos-Caro J., González G. A., 2008, MNRAS 390, 1587.

Pierens A., Hure J-M. 2004, ApJ, 605, 179.

Ramos-Caro J., López-Suspez F., González G. A., 2008, MNRAS 386, 440.

Schödel R. et al., 2002, Nature, 419, 694.

Verheinjen M. A. W., Sancici R., 2001, A&A, 370, 765.